



OUT- PATIENT AND PREVENTIVE HEALTH STATISTICAL BRIEF- 2019

March 2021

Private Bag 0024, Gaborone

Tel: 3671300 **Fax:** 3952201

Toll Free: 0800 600 200

Private Bag F193, City of Francistown

Tel. 241 5848, **Fax.** 241 7540

Private Bag 47, Maun

Tel: 371 5716 **Fax:** 686 4327

Private Bag 32 ,Ghanzi

Tel: 371 5723 **Fax:** 659 7506

E-mail: info@statsbots.org.bw

Website: <http://www.statsbots.org.bw>



STATISTICS BOTSWANA

PREFACE

This statistical brief focuses on analyzing the General Out-Patient and Preventive Health Statistics in Botswana. The information is derived from administrative data from patients seen at health facilities countrywide in 2019. It also shows the trends in Out-Patient statistics over a ten year period covering 2009 to 2019. The information provided will help in health policy formulation, monitoring and evaluation purposes.

The report shows that between 2018 and 2019 Out-Patient attendees on Notifiable Diseases increased from 451,537 in 2018 to 481,063 in 2019, Sexually Transmitted Infections (STI's) increased from 515,812 in 2018 to 551,694 in 2019. Ante Natal Care (ANC) increased from 367,489 in 2018 to 375,998 in 2019, Other Diseases increased from 3,667,421 in 2018 3,892,672 in 2019. Out-Patient attendees for other services, like Family Planning, Well Baby Care, Dressings and Tetanus Toxoid have also shown an increase. The report also shows that there was an outbreak of diarrhoea for week 51 only in 2019.

For more information and further enquiries, contact the Directorate of Stakeholder Relations on +267 367 1300. This publication, and all other Statistics Botswana outputs/publications are available on the website at ([http:// www.statsbots.org.bw](http://www.statsbots.org.bw)) and at the Statistics Botswana Information Resource Centre (Head-Office, Gaborone).

We sincerely thank all stakeholders involved in the formulation of this brief especially the Integrated Disease Surveillance (IDSR) in the Ministry of Health and Wellness for their continued support, as we strive to better serve users of our statistical products and services.



Dr Burton Mguni
Statistician General
March 2021

Table of Contents

| | |
|--|----|
| Preface..... | i |
| Table Of Contents..... | ii |
| 1.0 Introduction..... | 1 |
| 2.0 Summary Of Findings..... | 1 |
| 2.2 Sexually Transmitted Infections (S.T.I)..... | 1 |
| 2.3 Family Planning (Fp)-2019..... | 2 |
| 2.4 Antenatal Care..... | 3 |
| 2.5 Notifiable Diseases..... | 3 |
| Annexes: Tables..... | 6 |



1.0 INTRODUCTION

This statistical brief analyses administrative data from all health facilities in Botswana providing Out-Patient and Preventive Health Care. Out-Patient care is a medical procedure, tests and services that can be provided to patients in a setting that doesn't involve an overnight stay. All the 738 health facilities (35 Hospitals, 357 clinics, 346 health posts) and 1,117 mobile stops provide Out-Patient care in Botswana. Data is aggregated using the Out-Patient and Preventive Health Statistics Data Collection Tools which were implemented in response to the introduction of the International Classification of diseases 10th Edition (ICD) codes.

A consultation is either recorded as a new or repeat case. A new consultation refers to a case where a patient is seeking health services for a condition for the first time, whereas repeat refers to a case where a patient is getting health service for the same condition for the second time, or several times.

2.0 SUMMARY OF FINDINGS

2.1 Out-Patient Morbidity

Table 2.1.1 shows the top ten causes of morbidity in 2019. A total of 4,925,429 disease/conditions were recorded for Out-Patient attendances in 2019, which is an increase of 6.3 percent from 4,634,770 conditions recorded in 2018. New cases constituted 63.1 percent of all attendances. The highest cause of reported morbidity was Other diseases/conditions (34.0%), followed by Hypertension (12.5%), and Diseases of the Musculoskeletal System (11.6 %). These top three conditions were also identified in 2018. All diseases/conditions in the 2018 top ten list appear in the 2019 list though their ranks differ for some diseases.

Table 2.1.1 Top 10 Causes of Out-Patient Morbidity-2019

| Codes | Diagnosis | 2019 | | | 2018 | | |
|---------------------------------------|--|------|------------------|-------------|------|------------------|-------------|
| | | Rank | Total Attendance | Percent | Rank | Total Attendance | Percent |
| R68.8 | Other diseases/conditions | 1 | 1,675,094 | 34.0 | 1 | 1,617,131 | 34.9 |
| I10 | Hypertension | 2 | 613,940 | 12.5 | 2 | 566,411 | 12.2 |
| M79.9 | Diseases of the Musculoskeletal System | 3 | 573,805 | 11.6 | 3 | 521,813 | 11.3 |
| Z21 | HIV Positive | 4 | 357,831 | 7.3 | 6 | 265,391 | 5.7 |
| J11.1 | Influenza-like illness | 5 | 331,336 | 6.7 | 4 | 326,586 | 7.0 |
| L98.9 | Skin conditions | 6 | 271,550 | 5.5 | 5 | 306,120 | 6.6 |
| J03.9 | Tonsillitis | 7 | 227,809 | 4.6 | 7 | 199,928 | 4.3 |
| H57.9 | Eye diseases/conditions | 8 | 161,066 | 3.3 | 8 | 143,179 | 3.1 |
| X59.9 | Other external causes of injuries | 9 | 98,477 | 2.0 | 9 | 90,930 | 2.0 |
| K52.9 | Diarrhoea (acute) with no dehydration | 10 | 80,829 | 1.6 | 10 | 89,889 | 1.9 |
| Total (Causes specified above) | | | 4,391,737 | 89.2 | | 4,127,378 | 89.1 |
| Other Causes | | | 533,692 | 10.8 | | 507,392 | 10.9 |
| Total | | | 4,925,429 | 100 | | 4,634,770 | 100 |

2.2 Sexually Transmitted Infections (S.T.I)

A total of 551,694 sexually transmitted attendances were recorded in 2019. This is an increase of 7.0 percent from 515,812 cases recorded in 2018. Out of the 551,694 total attendances, 30.4 percent (167,745) were new attendances. **Figure 2.2.1** further shows an increase in the number of STI attendances from 2015 to 2016 followed by a slight decline of 7.1 percent in 2017, before registering an increase in 2018 and 2019.

Figure 2.2.1 Number of Sexually Transmitted Infections 2015-2019





Figure 2.2.2 shows distribution of STI attendance in 2019 by type. The highest recorded attendances were those who were HIV positive (64.9%) followed by those presenting Vaginal discharge Syndrome (12.8%), and Urethral discharge Syndrome (7.2%). The distribution of cases was higher among females (62.0%) compared to their male counterparts. Among the age groups, STI cases were higher within the age group 30 years and above (73.5%), and the lowest for age group 0-14 at 1.6 percent (Refer to Table 3.1c).


Figure 2.2.2 Sexually Transmitted Infections (STI's) Attendances -2019



2.3 Family Planning (FP)-2019

Family Planning allows individuals and couples to anticipate and attain their desired number of children and the spacing and timing of their births". It is achieved through use of contraceptive methods and the treatment of involuntary infertility. The Figure 2.3.1 below shows the distribution of family planning attendances by the preferred method of contraception. A total of 337,304 attendances were recorded in 2019, which is a decrease of 7.8 percent from 365,839 attendances recorded in 2018. New attendances were 57,256 (17.0%) compared to repeat attendances 280,048 (83.0%). The preferred method of contraception was male condom (56%) followed by Pill (22%) and the least being IUCD (1%).


Figure 2.3.1 Percentage Distribution of Family Planning Attendances-2019



2.4 Antenatal Care

Antenatal care is the care given to mothers from health professionals during pregnancy. It is sometimes called pregnancy care or maternity care. It is offered through appointments with a Registered Nurse midwife, or sometimes a doctor who specializes in pregnancy and birth (an Obstetrician). The total ANC visits for 2019 stood at 375,998 of which 52,330 (13.9%) were new attendances. This shows an increase of 2.3 Percent from 2018. The majority of new attendances were in the age group 30-39 (30.4%), followed by the age group 20-24 (26.6%) and age group 25-29 (26.3%). **Figure 2.4.1** shows a slightly fluctuating trend in ANC visits over the 5 year period. ANC attendances for those less than 20 years stood at 6,606 (13%) of the new attendances. **Figure 2.4.1** further reflects a constant trend in the percentage attendance for those less than 20 years except for a slight decrease noticed in 2017.

Figure 2.4.1: Number of ANC Visits 2015-2019




2.5 Notifiable Diseases

A notifiable disease is any disease that is required by law to be reported to government authorities. Malaria, Diarrhoea and Measles are some of communicable diseases that should be reported at the earliest time to the health authorities. Reporting of these diseases allows the authorities to monitor the diseases and provide for early warning of possible outbreaks. **Figure 2.5.1** shows number of confirmed

malaria and suspected measles cases between 2009 and 2019. Generally the figure reflects marginal fluctuations across the years for both malaria and suspected measles cases. A decrease is observed for malaria cases between the last three years, recording 239 malaria cases in 2019 compared to 553 cases in 2018. The number of suspected measles recorded a peak in 2010 (8,605 cases) followed by 4,205 and 3,441 cases in 2014 and 2011 respectively. The year 2019 has recorded the lowest cases (132) of suspected measles over the 2009-2019 interval.

Figure 2.5.1: Trends in Selected Notifiable Diseases, Botswana 2009-2019



A total of 8,489 diarrhea cases were reported in 2019, resulting in 55 deaths. This is a decrease of 35.7% from 13,208 cases registered in 2018. Generally, total number of reported diarrhea cases has been decreasing since 2009. An increase/decrease in diarrhea cases yields an increase/decrease in deaths. During the period 2011-2012 an inverse relation is observed where number of deaths increase over a decline in number of diarrhea cases. The same pattern was observed between 2018 and 2019.

Figure 2.5.2: Trends in Out-Patient Attendances and Deaths for Diarrhea, Botswana 2009-2019




Figure 2.5.3 shows the diarrhea case fatality over the ten-year period. Diarrhea Case fatality rate can be defined as the probability of death among cases diagnosed with diarrhea. The case fatality rate is often used as a measure of severity of illness. Diarrhea Case Fatality for 2019 was 6.5 deaths per 1,000 diarrhea cases; an increase from 3.9 deaths per 1,000 diarrhea cases recorded in 2018. Case fatality rates have been fluctuating over the years. The graph shows that the severity of diarrhea was highest in 2012 (9.3) and low in 2015 (4.0), 2016 (4.2), and 2018 (3.9) being the lowest.

Figure 2.5.3: Case Fatality Rate (Per 1,000) – 2019





Figure 2.5.4 shows the national diarrhoea thresholds for the year 2019. These thresholds serve as a reference point for health officials to be on alert and take action if there is an outbreak of diarrhoea in the country. In 2019, diarrhoea thresholds were below the alert and action threshold except for week 51 where thresholds were higher for alert and action. This shows that there was an outbreak of diarrhoea in week 51. The highest number of cases were recorded in week 41 (277 cases) and the lowest in week 29 (36 cases).

Figure 2.5.4: National Diarrhoea Thresholds- 2019



Annexes: Tables

Table 3.1a Number of Outpatient Attendances by Diagnosis, Age group and Sex (Notifiable Diseases)-2019

| Codes | Diagnosis | New Attendance | | | | | | | | Total |
|--------------|---------------------------------------|----------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|----------------|
| | | <1 | | 1-4 | | 5+ | | Sub Total | | |
| | | M | F | M | F | M | F | M | F | |
| A01.0 | Typhoid Fever | 19 | 870 | 45 | 1,193 | 18 | 1,033 | 82 | 3,096 | 3,178 |
| A09 | Diarrhoea with Blood | 581 | 489 | 800 | 690 | 753 | 809 | 2,134 | 1,988 | 4,122 |
| A36.9 | Diphtheria | 6 | 104 | 16 | 144 | 14 | 70 | 36 | 318 | 354 |
| A37.9 | Whooping Cough | 26 | 40 | 63 | 59 | 62 | 69 | 151 | 168 | 319 |
| B05.9 | Suspected Measles | 18 | 6 | 55 | 44 | 67 | 135 | 140 | 185 | 325 |
| B54 | Total confirmed malaria | 3 | 1 | 10 | 7 | 146 | 40 | 159 | 48 | 207 |
| B54.0 | Total Suspected Malaria | 597 | 473 | 1,751 | 1,519 | 5,239 | 6,073 | 7,587 | 8,065 | 15,652 |
| B54.1 | Total tested malaria | 518 | 402 | 1,655 | 1,295 | 5,369 | 5,538 | 7,542 | 7,235 | 14,777 |
| E43 | Severe acute malnutrition | 69 | 43 | 202 | 163 | 46 | 36 | 317 | 242 | 559 |
| E86 | Diarrhoea (acute) with dehydration | 1,050 | 869 | 1,330 | 1,258 | 1,316 | 1,478 | 3,696 | 3,605 | 7,301 |
| J11.1 | Influenza-like illness | 30,003 | 29,710 | 46,176 | 46,209 | 60,279 | 72,461 | 136,458 | 148,380 | 284,838 |
| J18.9 | Pneumonia | 1,313 | 2,508 | 1,675 | 3,849 | 1,007 | 3,852 | 3,995 | 10,209 | 14,204 |
| K52.9 | Diarrhoea (acute) with no dehydration | 9,912 | 8,700 | 14,345 | 12,961 | 12,751 | 14,849 | 37,008 | 36,510 | 73,518 |
| O98.6 | Total Confirmed Malaria in Pregnancy | . | 11 | . | 23 | . | 23 | - | 57 | 57 |
| Z20.3 | Rabies exposure | 46 | 20 | 147 | 98 | 1,383 | 968 | 1,576 | 1,086 | 2,662 |
| Total | | 44,161 | 44,246 | 68,270 | 69,512 | 88,450 | 107,434 | 200,881 | 221,192 | 422,073 |

Table 3.1a Number of Outpatient Attendances by Diagnosis, Age group and Sex (Notifiable Diseases)-2019
continued

| Codes | Diagnosis | Repeat Attendance | | | | | | | | Total | Grand Total |
|--------------|---------------------------------------|-------------------|--------------|--------------|--------------|---------------|---------------|---------------|---------------|---------------|----------------|
| | | <1 | | 1-4 | | 5+ | | Sub Total | | | |
| | | M | F | M | F | M | F | M | F | | |
| A01.0 | Typhoid Fever | 3 | 90 | 4 | 111 | 2 | 106 | 9 | 307 | 316 | 3,494 |
| A09 | Diarrhoea with Blood | 46 | 43 | 76 | 54 | 120 | 69 | 242 | 166 | 408 | 4,530 |
| A36.9 | Diphtheria | 1 | 21 | 2 | 14 | - | 10 | 3 | 45 | 48 | 402 |
| A37.9 | Whooping Cough | 8 | 13 | 39 | 46 | 57 | 66 | 104 | 125 | 229 | 548 |
| B05.9 | Suspected Measles | 1 | 7 | 2 | 3 | 4 | 4 | 7 | 14 | 21 | 346 |
| B54 | Total confirmed malaria | - | - | - | - | 4 | 4 | 4 | 4 | 8 | 215 |
| B54.0 | Total Suspected Malaria | 6 | 5 | 31 | 29 | 56 | 38 | 93 | 72 | 165 | 15,817 |
| B54.1 | Total tested malaria | 3 | 5 | 13 | 25 | 28 | 51 | 44 | 81 | 125 | 14,902 |
| E43 | Severe acute malnutrition | 16 | 14 | 220 | 219 | 41 | 45 | 277 | 278 | 555 | 1,114 |
| E86 | Diarrhoea (acute) with dehydration | 102 | 79 | 107 | 131 | 143 | 173 | 352 | 383 | 735 | 8,036 |
| J11.1 | Influenza-like illness | 4,375 | 4,764 | 6,891 | 6,990 | 10,787 | 12,691 | 22,053 | 24,445 | 46,498 | 331,336 |
| J18.9 | Pneumonia | 151 | 328 | 298 | 615 | 208 | 748 | 657 | 1,691 | 2,348 | 16,552 |
| K52.9 | Diarrhoea (acute) with no dehydration | 831 | 764 | 1,539 | 1,338 | 1,375 | 1,464 | 3,745 | 3,566 | 7,311 | 80,829 |
| O98.6 | Total Confirmed Malaria in Pregnancy | . | 3 | . | 19 | . | 6 | - | 28 | 28 | 85 |
| Z20.3 | Rabies exposure | 9 | 1 | 19 | 19 | 87 | 60 | 115 | 80 | 195 | 2,857 |
| Total | | 5,552 | 6,137 | 9,241 | 9,613 | 12,912 | 15,535 | 27,705 | 31,285 | 58,990 | 481,063 |

**Table 3.1b Number of Outpatient Attendances by Diagnosis, Age group and Sex
(Other Diseases/Conditions)-2019**

| Codes | Diagnosis | New Attendance | | | | | | | | Total |
|--------------|--|----------------|----------------|----------------|----------------|----------------|----------------|------------------|------------------|------------------|
| | | 0-14 | | 15-29 | | 30+ | | Sub total | | |
| | | M | F | M | F | M | F | M | F | |
| A16.9 | Tuberculosis | 83 | 98 | 211 | 225 | 836 | 513 | 1,130 | 836 | 1,966 |
| A30.9 | Leprosy | 146 | 144 | 39 | 47 | 48 | 69 | 233 | 260 | 493 |
| A64 | Stroke | 43 | 59 | 40 | 44 | 159 | 151 | 242 | 254 | 496 |
| B65.9 | Schistosomiasis | 166 | 173 | 59 | 64 | 50 | 79 | 275 | 316 | 591 |
| B83.9 | Helminthiasis | 807 | 935 | 343 | 496 | 396 | 568 | 1,546 | 1,999 | 3,545 |
| E10.8 | Diabetes Mellitus type 1 | 86 | 40 | 110 | 89 | 260 | 300 | 456 | 429 | 885 |
| E11.8 | Diabetes Mellitus type 2 | 57 | 74 | 109 | 190 | 604 | 944 | 770 | 1,208 | 1,978 |
| E46 | Malnutrition | 868 | 854 | 94 | 54 | 177 | 187 | 1,139 | 1,095 | 2,234 |
| F99 | Mental disorders | 475 | 408 | 989 | 1,167 | 1,768 | 2,576 | 3,232 | 4,151 | 7,383 |
| H57.9 | Eye diseases/conditions | 18,628 | 18,498 | 13,471 | 18,010 | 22,219 | 31,735 | 54,318 | 68,243 | 122,561 |
| H93.9 | Ear diseases/conditions | 11,451 | 12,357 | 7,911 | 9,795 | 9,349 | 11,823 | 28,711 | 33,975 | 62,686 |
| I10 | Hypertension | 43 | 125 | 1,049 | 1,458 | 7,235 | 10,058 | 8,327 | 11,641 | 19,968 |
| I50.9 | Cardiac Failure | 142 | 178 | 51 | 47 | 122 | 196 | 315 | 421 | 736 |
| J03.9 | Tonsillitis | 37,293 | 46,359 | 20,501 | 33,133 | 19,738 | 34,999 | 77,532 | 114,491 | 192,023 |
| J45.9 | Asthma | 1,265 | 857 | 1,189 | 1,213 | 1,713 | 1,433 | 4,167 | 3,503 | 7,670 |
| L98.9 | Skin conditions | 51,078 | 51,356 | 24,093 | 31,530 | 29,735 | 41,058 | 104,906 | 123,944 | 228,850 |
| M79.9 | Diseases of the Musculoskeletal System | 10,524 | 11,842 | 41,229 | 50,359 | 112,429 | 182,403 | 164,182 | 244,604 | 408,786 |
| R68.8 | Other diseases/conditions | 192,647 | 206,758 | 135,630 | 223,513 | 225,786 | 370,176 | 554,063 | 800,447 | 1,354,510 |
| U88 | Suspected MDR-TB | 122 | 165 | 22 | 35 | 60 | 63 | 204 | 263 | 467 |
| U88.1 | Laboratory confirmed MDR-TB | 273 | 319 | 174 | 223 | 162 | 227 | 609 | 769 | 1,378 |
| U88.2 | Laboratory confirmed XDR-TB | 61 | 77 | 7 | 13 | 15 | 25 | 83 | 115 | 198 |
| V89.2 | Injuries secondary to road traffic accidents | 1,374 | 931 | 1,908 | 1,512 | 3,154 | 2,401 | 6,436 | 4,844 | 11,280 |
| X59.9 | Other external causes of injuries | 14,933 | 10,086 | 15,895 | 11,252 | 19,344 | 15,539 | 50,172 | 36,877 | 87,049 |
| Total | | 342,565 | 362,693 | 265,124 | 384,469 | 455,359 | 707,523 | 1,063,048 | 1,454,685 | 2,517,733 |

**Table 3.1b Number of Outpatient Attendances by Diagnosis, Age group and Sex
(Other Diseases/Conditions)-2019, continued**

| Codes | Diagnosis | Repeat Attendance | | | | | | | | Total | Grand Total |
|--------------|--|-------------------|---------------|---------------|----------------|----------------|----------------|----------------|----------------|------------------|------------------|
| | | 0-14 | | 15-29 | | 30+ | | Sub total | | | |
| | | M | F | M | F | M | F | M | F | | |
| A16.9 | Tuberculosis | 105 | 86 | 340 | 413 | 1,588 | 1,371 | 2,033 | 1,870 | 3,903 | 5,869 |
| A30.9 | Leprosy | 21 | 20 | 14 | 6 | 35 | 17 | 70 | 43 | 113 | 606 |
| A64 | Stroke | 3 | 1 | 32 | 35 | 2,203 | 751 | 2,238 | 787 | 3,025 | 3,521 |
| B65.9 | Schistosomiasis | 4 | 7 | 14 | 19 | 16 | 24 | 34 | 50 | 84 | 675 |
| B83.9 | Helminthiasis | 136 | 169 | 80 | 148 | 152 | 192 | 368 | 509 | 877 | 4,422 |
| E10.8 | Diabetes Mellitus type 1 | 86 | 70 | 455 | 618 | 3,612 | 6,232 | 4,153 | 6,920 | 11,073 | 11,958 |
| E11.8 | Diabetes Mellitus type 2 | 93 | 180 | 603 | 737 | 11,897 | 26,173 | 12,593 | 27,090 | 39,683 | 41,661 |
| E46 | Malnutrition | 2,101 | 2,244 | 85 | 162 | 1,191 | 3,593 | 3,377 | 5,999 | 9,376 | 11,610 |
| F99 | Mental disorders | 749 | 621 | 4,856 | 3,698 | 16,806 | 17,333 | 22,411 | 21,652 | 44,063 | 51,446 |
| H57.9 | Eye diseases/ conditions | 2,780 | 2,729 | 3,533 | 4,311 | 9,908 | 15,244 | 16,221 | 22,284 | 38,505 | 161,066 |
| H93.9 | Ear diseases/ conditions | 1,973 | 2,102 | 2,071 | 2,307 | 2,963 | 3,554 | 7,007 | 7,963 | 14,970 | 77,656 |
| I10 | Hypertension | 437 | 1,017 | 8,490 | 13,753 | 174,779 | 395,496 | 183,706 | 410,266 | 593,972 | 613,940 |
| I50.9 | Cardiac Failure | 12 | 19 | 28 | 22 | 684 | 1,104 | 724 | 1,145 | 1,869 | 2,605 |
| J03.9 | Tonsillitis | 5,188 | 6,709 | 4,168 | 6,400 | 5,022 | 8,299 | 14,378 | 21,408 | 35,786 | 227,809 |
| J45.9 | Asthma | 3,068 | 2,440 | 4,208 | 5,114 | 8,847 | 11,248 | 16,123 | 18,802 | 34,925 | 42,595 |
| L98.9 | Skin conditions | 6,552 | 6,655 | 4,887 | 6,028 | 7,663 | 10,915 | 19,102 | 23,598 | 42,700 | 271,550 |
| M79.9 | Diseases of the Musculoskeletal System | 3,285 | 4,390 | 12,595 | 16,021 | 45,544 | 83,184 | 61,424 | 103,595 | 165,019 | 573,805 |
| R68.8 | Other diseases/ conditions | 29,742 | 34,196 | 29,756 | 48,276 | 67,364 | 111,250 | 126,862 | 193,722 | 320,584 | 1,675,094 |
| U88 | Suspected MDR-TB | 24 | 29 | 10 | 9 | 26 | 31 | 60 | 69 | 129 | 596 |
| U88.1 | Laboratory confirmed MDR-TB | 40 | 64 | 49 | 73 | 55 | 63 | 144 | 200 | 344 | 1,722 |
| U88.2 | Laboratory confirmed XDR-TB | 36 | 46 | 2 | 3 | 32 | 65 | 70 | 114 | 184 | 382 |
| V89.2 | Injuries secondary to road traffic accidents | 185 | 126 | 301 | 285 | 829 | 601 | 1,315 | 1,012 | 2,327 | 13,607 |
| X59.9 | Other external causes of injuries | 1,349 | 1,034 | 1,855 | 1,506 | 2,882 | 2,802 | 6,086 | 5,342 | 11,428 | 98,477 |
| Total | | 57,969 | 64,954 | 78,432 | 109,944 | 364,098 | 699,542 | 500,499 | 874,440 | 1,374,939 | 3,892,672 |

**Table 3.1c Number of Outpatient Attendances by Diagnosis, Age group and Sex
(Sexually Transmitted Infections)-2019**

| Codes | Diagnosis | New Attendance | | | | | | | | Total |
|--------------|--------------------------------|----------------|--------------|---------------|---------------|---------------|---------------|---------------|----------------|----------------|
| | | 0-14 | | 15-29 | | 30+ | | Sub Total | | |
| | | M | F | M | F | M | F | M | F | |
| A53.9 | Syphilis RPT+/VDRL+) | 32 | 12 | 311 | 422 | 473 | 443 | 816 | 877 | 1,693 |
| A63.0 | Genital Warts | 15 | 25 | 316 | 423 | 339 | 405 | 670 | 853 | 1,523 |
| A64 | Other STI. (e.g Inguinal Bubo) | 6 | 14 | 271 | 213 | 325 | 259 | 602 | 486 | 1,088 |
| B16.9 | Hepatitis B | 2 | 1 | 3 | 1 | 11 | 10 | 16 | 12 | 28 |
| B24 | AIDS | 2 | 5 | 44 | 64 | 250 | 166 | 296 | 235 | 531 |
| N34.3 | Urethral discharge Syndrome | 348 | . | 16,150 | . | 18,852 | . | 35,350 | - | 35,350 |
| N50.9 | Genital ulcer syndrome | 107 | . | 2,716 | . | 3,642 | . | 6,465 | - | 6,465 |
| N73.9 | Lower abdominal pains (PID) | . | 191 | . | 9,422 | . | 11,362 | - | 20,975 | 20,975 |
| N76.6 | Genital ulcer syndrome | . | 75 | . | 3,503 | . | 3,682 | - | 7,260 | 7,260 |
| N89.8 | Vaginal discharge Syndrome | . | 765 | . | 32,097 | . | 28,395 | - | 61,257 | 61,257 |
| Z20.2 | STI Contacts | 208 | 219 | 5,296 | 4,933 | 7,527 | 5,479 | 13,031 | 10,631 | 23,662 |
| Z21 | HIV Positive | 91 | 132 | 821 | 1,419 | 2,454 | 2,996 | 3,366 | 4,547 | 7,913 |
| Total | | 811 | 1,439 | 25,928 | 52,497 | 33,873 | 53,197 | 60,612 | 107,133 | 167,745 |

**Table 3.1c Number of Outpatient Attendances by Diagnosis, Age group and Sex
(Sexually Transmitted Infections)-2019, continued**

| Codes | Diagnosis | Repeat Attendance | | | | | | | | Total | Grand Total |
|--------------|--------------------------------|-------------------|--------------|---------------|---------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | | 0-14 | | 15-29 | | 30+ | | Sub Total | | | |
| | | M | F | M | F | M | F | M | F | | |
| A53.9 | Syphilis RPT+/VDRL+) | 3 | 4 | 24 | 44 | 76 | 56 | 103 | 104 | 207 | 1,900 |
| A63.0 | Genital Warts | 5 | 2 | 56 | 89 | 91 | 150 | 152 | 241 | 393 | 1,916 |
| A64 | Other STI. (e.g Inguinal Bubo) | 4 | - | 34 | 17 | 47 | 36 | 85 | 53 | 138 | 1,226 |
| B16.9 | Hepatitis B | 4 | 7 | 20 | 12 | 82 | 179 | 106 | 198 | 304 | 332 |
| B24 | AIDS | 81 | 84 | 643 | 652 | 4,878 | 5,097 | 5,602 | 5,833 | 11,435 | 11,966 |
| N34.3 | Urethral discharge Syndrome | 53 | . | 1,848 | . | 2,391 | . | 4,292 | - | 4,292 | 39,642 |
| N50.9 | Genital ulcer syndrome | 13 | . | 398 | . | 598 | . | 1,009 | - | 1,009 | 7,474 |
| N73.9 | Lower abdominal pains (PID) | . | 45 | . | 1,561 | . | 1,991 | - | 3,597 | 3,597 | 24,572 |
| N76.6 | Genital ulcer syndrome | . | 8 | . | 472 | . | 633 | - | 1,113 | 1,113 | 8,373 |
| N89.8 | Vaginal discharge Syndrome | . | 120 | . | 4,340 | . | 4,721 | - | 9,181 | 9,181 | 70,438 |
| Z20.2 | STI Contacts | 15 | 9 | 513 | 498 | 696 | 631 | 1,224 | 1,138 | 2,362 | 26,024 |
| Z21 | HIV Positive | 2,855 | 3,472 | 16,930 | 30,791 | 116,489 | 179,381 | 136,274 | 213,644 | 349,918 | 357,831 |
| Total | | 3,033 | 3,751 | 20,466 | 38,476 | 125,348 | 192,875 | 148,847 | 235,102 | 383,949 | 551,694 |

**Table 3.2 Number of Outpatient Attendances by Health District and Diagnosis
(Complicated Pregnancies)-2019**

| Health District | Anaemia <10g/l | Anaemia ≥10g/l | Haemorrhage | Infection | Hypertension |
|-----------------|----------------|----------------|-------------|--------------|---------------|
| Ngamiland | 860 | 614 | 214 | 272 | 542 |
| North East | 294 | 290 | 8 | 52 | 136 |
| Palapye | 672 | 672 | 14 | 180 | 416 |
| Bobirwa | 600 | 1,418 | 8 | 40 | 222 |
| Kweneng East | 504 | 1,096 | 30 | 308 | 1,170 |
| Southern | 494 | 298 | 16 | 62 | 292 |
| Gantsi | 102 | 104 | 2 | 8 | 124 |
| Mahalapye | 892 | 640 | 48 | 288 | 694 |
| Kgatleng | 550 | 704 | 12 | 104 | 822 |
| Chobe | 660 | 1,986 | 4 | 104 | 138 |
| Kgalagadi South | 86 | 108 | 28 | 58 | 244 |
| Tutume | 796 | 1,082 | 18 | 160 | 188 |
| Boteti | 360 | 142 | 6 | 126 | 164 |
| Okavango | 422 | 250 | 20 | 42 | 936 |
| Gaborone | 4,076 | 12,776 | 148 | 3,580 | 2,410 |
| Francistown | 2,644 | 3,300 | 98 | 826 | 1,074 |
| South East | 504 | 250 | 20 | 198 | 380 |
| Lobatse | 466 | 164 | 30 | 202 | 926 |
| Selibe Phikwe | 308 | 274 | 2 | 114 | 134 |
| Kweneng West | 90 | 122 | 110 | 30 | 118 |
| Mabutsane | 62 | 504 | - | - | 14 |
| Jwaneng | 316 | 402 | 4 | 104 | 108 |
| Goodhope | 186 | 216 | 34 | 224 | 280 |
| Kgalagadi North | 80 | 166 | - | 52 | 60 |
| Moshupa | 100 | 86 | - | 36 | 86 |
| Charleshill | 38 | 4 | - | 34 | 36 |
| Serowe | 1,072 | 2,042 | 30 | 496 | 788 |
| Total | 17,234 | 29,710 | 904 | 7,700 | 12,502 |

**Table 3.3 Number of Outpatient Attendances by Health District and Diagnosis
(Other Services)-2019**

| Health district | Well Baby Care | | Injection | Dressings | Tetanus Toxoid |
|-----------------|----------------|------------------|----------------|----------------|----------------|
| | New | Repeat | | | |
| Ngamiland | 2,603 | 93,734 | 17,860 | 9,540 | 1,370 |
| North East | 1,075 | 53,969 | 6,286 | 5,649 | 569 |
| Palapye | 2,122 | 110,749 | 17,160 | 14,562 | 1,145 |
| Bobirwa | 2,116 | 81,533 | 12,121 | 9,533 | 1,108 |
| Kweneng East | 3,723 | 152,854 | 21,014 | 21,193 | 1,627 |
| Southern | 1,457 | 72,728 | 9,939 | 7,610 | 691 |
| Gantsi | 553 | 36,828 | 4,394 | 4,008 | 644 |
| Mahalapye | 4,679 | 146,425 | 17,600 | 20,024 | 1,708 |
| Kgatleng | 1,682 | 110,587 | 18,202 | 17,536 | 1,320 |
| Chobe | 302 | 21,475 | 5,461 | 4,673 | 587 |
| Kgalagadi South | 868 | 36,487 | 4,327 | 2,412 | 433 |
| Tutume | 5,586 | 94,366 | 15,660 | 10,253 | 1,087 |
| Boteti | 2,304 | 84,658 | 9,696 | 7,088 | 870 |
| Okavango | 2,140 | 63,786 | 7,877 | 4,619 | 1,426 |
| Gaborone | 7,702 | 248,320 | 97,269 | 65,184 | 5,077 |
| Francistown | 2,122 | 171,344 | 41,660 | 31,639 | 2,301 |
| South East | 725 | 43,799 | 9,366 | 10,362 | 1,021 |
| Lobatse | 1,582 | 37,819 | 10,616 | 9,017 | 622 |
| Selibe Phikwe | 1,131 | 44,655 | 13,496 | 8,891 | 246 |
| Kweneng West | 2,170 | 53,916 | 6,193 | 6,506 | 1,278 |
| Mabutsane | 395 | 18,685 | 1,812 | 1,503 | 277 |
| Jwaneng | 503 | 31,451 | 5,775 | 3,698 | 542 |
| Goodhope | 1,766 | 50,385 | 7,024 | 5,427 | 697 |
| Kgalagadi North | 392 | 22,779 | 3,833 | 2,664 | 307 |
| Moshupa | 1,084 | 51,660 | 8,599 | 6,576 | 491 |
| Charleshill | 224 | 13,963 | 2,266 | 2,565 | 372 |
| Serowe | 2,092 | 118,258 | 20,407 | 18,197 | 1,580 |
| Total | 53,098 | 2,067,213 | 395,913 | 310,929 | 29,396 |

Table 3.4 Number of Out-patient attendances by Health district and Diagnosis (Family Planning) - 2019

| Health district | New attendance | | | | | | | | | | Total |
|-----------------|----------------|----------------|-------------------|--------------|-------------|--------------|---------------|-----------------|--------------|------------|---------------|
| | Pill Low Dose | Pill High Dose | Pill Progesterone | IUCDS Copper | IUCDS Other | Injections | Condom (Male) | Condom (Female) | Nor-Plant | Other | |
| Ngamiland | 413 | 42 | 175 | 1 | - | 224 | 318 | 20 | 94 | 26 | 1,313 |
| North East | 111 | 14 | 11 | 1 | - | 114 | 62 | 7 | 67 | 4 | 391 |
| Palapye | 269 | 72 | 157 | 9 | - | 368 | 970 | 42 | 256 | 14 | 2,157 |
| Bobirwa | 363 | 76 | 247 | 3 | - | 215 | 640 | 53 | 131 | 21 | 1,749 |
| Kweneng East | 664 | 51 | 332 | 13 | 2 | 389 | 1,331 | 62 | 77 | 46 | 2,967 |
| Southern | 147 | 44 | 84 | - | - | 207 | 1,509 | 125 | 112 | 8 | 2,236 |
| Gantsi | 107 | 17 | 96 | - | - | 202 | 559 | 15 | 155 | 21 | 1,172 |
| Mahalapye | 807 | 104 | 416 | 45 | 4 | 603 | 1,390 | 128 | 249 | 35 | 3,781 |
| Kgatleng | 532 | 73 | 190 | 13 | 4 | 461 | 2,094 | 107 | 315 | 29 | 3,818 |
| Chobe | 231 | 1 | 42 | 10 | - | 134 | 6,940 | 45 | 41 | 1 | 7,445 |
| Kgalagadi South | 66 | 64 | 100 | - | 7 | 104 | 177 | 27 | 47 | 17 | 609 |
| Tutume | 282 | 44 | 168 | 4 | 2 | 295 | 1,321 | 87 | 101 | 6 | 2,310 |
| Boteti | 469 | 101 | 102 | 10 | - | 446 | 625 | 88 | 236 | 31 | 2,108 |
| Okavango | 308 | 40 | 158 | - | - | 94 | 3,170 | 37 | 32 | 3 | 3,842 |
| Gaborone | 780 | 55 | 477 | 142 | 9 | 846 | 5,451 | 104 | 406 | 43 | 8,313 |
| Francistown | 661 | 46 | 276 | 22 | 30 | 341 | - | 2 | 374 | 37 | 1,789 |
| South East | 148 | - | 115 | 37 | 4 | 110 | 2 | - | 19 | 69 | 504 |
| Lobatse | 170 | 10 | 66 | 13 | - | 76 | 548 | 133 | 217 | 5 | 1,238 |
| Selibe Phikwe | 112 | 2 | 74 | 8 | - | 162 | 200 | - | 71 | 6 | 635 |
| Kweneng West | 247 | 6 | 233 | 8 | 2 | 59 | 1,263 | 15 | 131 | 22 | 1,986 |
| Mabutsane | 97 | 20 | 52 | - | - | 81 | 193 | 1 | 3 | 9 | 456 |
| Jwaneng | 55 | 1 | 14 | 5 | 1 | 82 | 1,037 | 1 | 34 | 6 | 1,236 |
| Goodhope | 260 | 54 | 60 | - | - | 292 | 692 | 24 | 59 | 12 | 1,453 |
| Kgalagadi North | 80 | - | 33 | 3 | 1 | 90 | 211 | 55 | 16 | - | 489 |
| Moshupa | 101 | 14 | 65 | 10 | - | 247 | 1,297 | 68 | 65 | 19 | 1,886 |
| Charleshill | 71 | 58 | 58 | 2 | - | 116 | - | - | 26 | 4 | 335 |
| Serowe | 275 | 18 | 159 | 20 | - | 222 | 49 | 30 | - | 265 | 1,038 |
| Total | 7,826 | 1,027 | 3,960 | 379 | 66 | 6,580 | 32,049 | 1,276 | 3,334 | 759 | 57,256 |

Table 3.5 Number of Out-patient attendances by Health district and Diagnosis (Family Planning) - 2019, continued


| Health district | Repeat attendance | | | | | | | | | | Total |
|-----------------|-------------------|----------------|-------------------|--------------|-------------|---------------|----------------|-----------------|--------------|--------------|----------------|
| | Pill Low Dose | Pill High Dose | Pill Progesterone | IUCDS Copper | IUCDS Other | Injections | Condom (Male) | Condom (Female) | Nor-Plant | Other | |
| Ngamiland | 3,344 | 233 | 1,070 | 10 | - | 1,179 | 5,726 | 45 | 180 | 119 | 11,906 |
| North East | 1,564 | 35 | 133 | 13 | 14 | 1,524 | 316 | 19 | 185 | 120 | 3,923 |
| Palapye | 2,336 | 304 | 1,064 | 31 | - | 3,550 | 7,401 | 94 | 445 | 49 | 15,274 |
| Bobirwa | 2,345 | 275 | 1,026 | 19 | 5 | 1,331 | 3,978 | 247 | 104 | 1 | 9,331 |
| Kweneng East | 5,116 | 342 | 1,550 | 24 | 2 | 1,618 | 5,065 | 65 | 90 | 30 | 13,902 |
| Southern | 1,207 | 206 | 997 | 12 | 11 | 1,509 | 6,243 | 244 | 142 | 50 | 10,621 |
| Gantsi | 686 | 184 | 829 | 4 | 1 | 939 | 1,269 | 2 | 109 | 23 | 4,046 |
| Mahalapye | 4,682 | 448 | 1,285 | 45 | 1 | 3,073 | 6,525 | 406 | 170 | 19 | 16,654 |
| Kgatleng | 4,334 | 266 | 822 | 26 | 2 | 1,313 | 6,126 | 106 | 211 | 39 | 13,245 |
| Chobe | 1,771 | 33 | 203 | 13 | - | 646 | 1,498 | 66 | 85 | 9 | 4,324 |
| Kgalagadi South | 920 | 312 | 713 | 1 | - | 521 | 1,461 | 37 | 12 | 19 | 3,996 |
| Tutume | 3,061 | 338 | 936 | 44 | - | 2,611 | 12,055 | 3,175 | 294 | 69 | 22,583 |
| Boteti | 2,852 | 347 | 585 | 30 | 2 | 2,726 | 6,252 | 420 | 162 | 52 | 13,428 |
| Okavango | 3,236 | 419 | 2,888 | - | 1 | 2,096 | 6,848 | 254 | 117 | 4 | 15,863 |
| Gaborone | 6,086 | 223 | 2,831 | 243 | 25 | 5,564 | 31,703 | 394 | 507 | 380 | 47,956 |
| Francistown | 5,096 | 212 | 1,129 | 80 | 63 | 2,171 | 847 | 3 | 1,031 | 165 | 10,797 |
| South East | 1,444 | 20 | 1,014 | 41 | 8 | 1,153 | 276 | - | 26 | 189 | 4,171 |
| Lobatse | 1,660 | 140 | 576 | 51 | - | 712 | 1,835 | 8 | 338 | 59 | 5,379 |
| Selibe Phikwe | 978 | 14 | 576 | 40 | - | 1,492 | 1,008 | 14 | 257 | 63 | 4,442 |
| Kweneng West | 2,373 | 150 | 1,773 | 21 | 8 | 312 | 3,226 | 79 | 163 | 101 | 8,206 |
| Mabutsane | 677 | 110 | 270 | - | - | 457 | 890 | 42 | 19 | 13 | 2,478 |
| Jwaneng | 1,513 | 2 | 432 | 19 | 6 | 1,504 | 2,507 | 24 | 64 | 59 | 6,130 |
| Goodhope | 2,058 | 246 | 273 | 31 | 15 | 2,042 | 3,763 | 167 | 92 | 79 | 8,766 |
| Kgalagadi North | 913 | 67 | 242 | 12 | - | 651 | 918 | 117 | 106 | 36 | 3,062 |
| Moshupa | 1,343 | 172 | 501 | 20 | 2 | 1,964 | 5,027 | 66 | 161 | 31 | 9,287 |
| Charleshill | 601 | 242 | 197 | - | - | 454 | - | 4 | 36 | 2 | 1,536 |
| Serowe | 3,083 | 223 | 1,370 | 20 | 1 | 2,282 | 1,197 | 52 | - | 514 | 8,742 |
| Total | 65,279 | 5,563 | 25,285 | 850 | 167 | 45,394 | 123,960 | 6,150 | 5,106 | 2,294 | 280,048 |

**Table 3.6 Number of Outpatient Attendances by Health District and Age group
(Ante Natal and Post Natal Care)-2019**

| District | New attendance | | | | | | | Total | ANC First Trimester | Post Natal care |
|-----------------|----------------|--------------|---------------|---------------|---------------|--------------|------------|---------------|---------------------|-----------------|
| | <15 | 15-19 | 20-24 | 25-29 | 30-39 | 40-44 | 45+ | | | |
| Ngamiland | 12 | 417 | 892 | 789 | 973 | 149 | 3 | 3,235 | 425 | 1,441 |
| North East | - | 89 | 202 | 178 | 226 | 26 | 1 | 722 | 178 | 844 |
| Palapye | 4 | 283 | 605 | 528 | 674 | 78 | 6 | 2,178 | 578 | 1,364 |
| Bobirwa | 3 | 195 | 376 | 333 | 353 | 40 | 10 | 1,310 | 181 | 1,135 |
| Kweneng East | 8 | 344 | 771 | 636 | 931 | 136 | 18 | 2,844 | 401 | 1,853 |
| Southern | 2 | 186 | 332 | 314 | 405 | 57 | 3 | 1,299 | 191 | 928 |
| Gantsi | 1 | 222 | 322 | 315 | 279 | 43 | 5 | 1,187 | 175 | 322 |
| Mahalapye | 6 | 384 | 646 | 540 | 782 | 90 | 5 | 2,453 | 451 | 1,888 |
| Kgatleng | 5 | 248 | 645 | 654 | 738 | 85 | 3 | 2,378 | 310 | 1,723 |
| Chobe | - | 84 | 228 | 415 | 270 | 28 | - | 1,025 | 272 | 357 |
| Kgalagadi South | 2 | 78 | 149 | 98 | 221 | 35 | - | 583 | 75 | 367 |
| Tutume | 16 | 325 | 425 | 442 | 452 | 66 | 5 | 1,731 | 300 | 1,176 |
| Boteti | 9 | 326 | 573 | 742 | 666 | 60 | 4 | 2,380 | 290 | 1,030 |
| Okavango | 19 | 274 | 500 | 460 | 580 | 87 | 10 | 1,930 | 145 | 1,045 |
| Gaborone | 7 | 872 | 3,274 | 3,382 | 3,398 | 365 | 15 | 11,313 | 2,865 | 4,112 |
| Francistown | 10 | 611 | 1,377 | 1,340 | 1,657 | 167 | 11 | 5,173 | 1,132 | 3,218 |
| South East | 1 | 112 | 248 | 328 | 435 | 60 | 1 | 1,185 | 284 | 896 |
| Lobatse | 1 | 115 | 279 | 269 | 377 | 37 | 1 | 1,079 | 151 | 741 |
| Selibe Phikwe | - | 115 | 256 | 266 | 368 | 53 | 5 | 1,063 | 258 | 657 |
| Kweneng West | 2 | 229 | 301 | 283 | 314 | 55 | 8 | 1,192 | 95 | 445 |
| Mabutsane | - | 54 | 86 | 74 | 90 | 14 | 1 | 319 | 52 | 198 |
| Jwaneng | - | 126 | 220 | 256 | 283 | 23 | 3 | 911 | 185 | 399 |
| Goodhope | 2 | 169 | 208 | 181 | 244 | 42 | 9 | 855 | 174 | 726 |
| Kgalagadi North | 1 | 86 | 125 | 140 | 160 | 21 | 6 | 539 | 118 | 282 |
| Moshupa | 5 | 125 | 200 | 162 | 238 | 30 | 7 | 767 | 106 | 773 |
| Charleshill | 2 | 76 | 99 | 119 | 104 | 19 | - | 419 | 44 | 157 |
| Serowe | 6 | 337 | 580 | 541 | 701 | 86 | 9 | 2,260 | 598 | 1,675 |
| Total | 124 | 6,482 | 13,919 | 13,785 | 15,919 | 1,952 | 149 | 52,330 | 10,034 | 29,752 |

**Table 3.7 Number of Outpatient Attendances by Health Districts and Age group
(Ante Natal and Post Natal Care)-2019, continued**

| District | Repeat attendance | | | | | | | Total | Total ANC Attendances |
|--------------------|-------------------|---------------|---------------|---------------|----------------|---------------|------------|----------------|-----------------------|
| | <15 | 15-19 | 20-24 | 25-29 | 30-39 | 40-44 | 45+ | | |
| Ngamiland | 44 | 1,450 | 3,737 | 3,787 | 4,623 | 494 | 32 | 14,167 | 17,402 |
| North East | 11 | 460 | 1,546 | 1,252 | 1,649 | 203 | 4 | 5,125 | 5,847 |
| Palapye | 16 | 1,318 | 3,693 | 3,556 | 4,746 | 597 | 17 | 13,943 | 16,121 |
| Bobirwa | 12 | 1,187 | 2,615 | 2,339 | 2,577 | 270 | 39 | 9,039 | 10,349 |
| Kweneng East | 54 | 1,916 | 4,846 | 4,817 | 5,796 | 814 | 37 | 18,280 | 21,124 |
| Southern | 9 | 957 | 2,397 | 2,520 | 2,779 | 298 | 7 | 8,967 | 10,266 |
| Gantsi | 8 | 935 | 1,480 | 1,605 | 1,491 | 175 | 14 | 5,708 | 6,895 |
| Mahalapye | 29 | 2,082 | 4,589 | 4,355 | 5,709 | 475 | 24 | 17,263 | 19,716 |
| Kgatleng | 22 | 1,340 | 4,232 | 4,221 | 4,931 | 510 | 25 | 15,281 | 17,659 |
| Chobe | - | 427 | 1,233 | 1,487 | 1,693 | 233 | 9 | 5,082 | 6,107 |
| Kgalagadi South | 9 | 415 | 767 | 565 | 1,379 | 175 | 1 | 3,311 | 3,894 |
| Tutume | 34 | 1,323 | 2,572 | 2,685 | 2,863 | 369 | 17 | 9,863 | 11,594 |
| Boteti | 29 | 1,410 | 3,233 | 3,342 | 4,316 | 372 | 26 | 12,728 | 15,108 |
| Okavango | 41 | 1,021 | 2,375 | 2,487 | 2,972 | 370 | 41 | 9,307 | 11,237 |
| Gaborone | 27 | 4,060 | 18,729 | 21,579 | 21,162 | 2,202 | 75 | 67,834 | 79,147 |
| Francistown | 35 | 3,269 | 9,732 | 9,827 | 12,142 | 1,342 | 67 | 36,414 | 41,587 |
| South East | 8 | 716 | 1,984 | 2,562 | 3,752 | 385 | 5 | 9,412 | 10,597 |
| Lobatse | 12 | 682 | 1,894 | 2,448 | 2,900 | 332 | 7 | 8,275 | 9,354 |
| Selibe Phikwe | 19 | 713 | 1,867 | 2,262 | 2,955 | 391 | 19 | 8,226 | 9,289 |
| Kweneng West | 4 | 1,014 | 1,630 | 1,867 | 1,969 | 245 | 33 | 6,762 | 7,954 |
| Mabutsane | - | 246 | 499 | 490 | 437 | 70 | 3 | 1,745 | 2,064 |
| Jwaneng | 16 | 615 | 1,501 | 1,936 | 2,044 | 146 | 23 | 6,281 | 7,192 |
| Goodhope | 21 | 795 | 1,495 | 1,348 | 1,568 | 208 | 18 | 5,453 | 6,308 |
| Kgalagadi North | 9 | 377 | 822 | 861 | 1,036 | 144 | 5 | 3,254 | 3,793 |
| Moshupa | 39 | 602 | 1,343 | 1,379 | 1,498 | 173 | 15 | 5,049 | 5,816 |
| Charleshill | 11 | 302 | 513 | 644 | 601 | 86 | - | 2,157 | 2,576 |
| Serowe | 41 | 1,867 | 3,957 | 3,694 | 4,521 | 617 | 45 | 14,742 | 17,002 |
| Grand total | 560 | 31,499 | 85,281 | 89,915 | 104,109 | 11,696 | 608 | 323,668 | 375,998 |



Private Bag 0024, Gaborone
Tel: 3671300 **Fax:** 3952201
Toll Free: 0800 600 200

Private Bag F193, City of Francistown
Tel. 241 5848, **Fax.** 241 7540

Private Bag 47, Maun
Tel: 371 5716 **Fax:** 686 4327

Private Bag 32 ,Ghanzi
Tel: 371 5723 **Fax:** 659 7506

E-mail: info@statsbots.org.bw

Website: <http://www.statsbots.org.bw>



STATISTICS BOTSWANA